arXiv:1505.03627v1 [math.DG] 14 May 2015

Killing Vector Fields on Multiply Warped Products with a Semi-symmetric Metric Connection

Quan Qu *

Abstract

In this paper, we define a semi-symmetric metric Killing vector field, then study semi-symmetric metric Killing vector fields on warped and multiply warped products with a semi-symmetric metric connection. We also study Killing and 2-Killing vector fields on multiply warped products.

Keywords: Warped products; multiply warped products; semi-symmetric metric connection; Killing vector fields; semi-symmetric metric Killing vector fields; 2-Killing vector fields.

2010 Mathematics Subject Classification. 53B05, 53C21, 53C25, 53C50, 53C80.

1 Introduction

The warped product $M_1 \times_f M_2$ of two pseudo-Riemannian manifolds (M_1, g_1) and (M_2, g_2) with a smooth function $f : M_1 \to (0, \infty)$ is a product manifold of form $M_1 \times M_2$ with the metric tensor $g = g_1 \oplus f^2 g_2$. Here, (M_1, g_1) is called the base manifold and (M_2, g_2) is called as the fiber manifold and f is called as the warping function. The concept of warped products was first introduced by Bishop and O'Neill in [1] to construct examples of Riemannian manifolds with negative curvature.

One can generalize warped products to multiply warped products. A multiply warped product (M, g) is the product manifold $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ with the metric $g = g_B \oplus f_1^2 g_{M_1} \oplus f_2^2 g_{M_2} \cdots \oplus f_m^2 g_{M_m}$, where for each $i \in \{1, \cdots, m\}$, $f_i : B \to (0, \infty)$ is smooth and (M_i, g_{M_i}) is a pseudo-Riemannian manifold.

Killing vector fields have been studied on Riemannian and pseudo-Riemannian manifolds for a long time. The problems of existence and characterization of Killing vector fields are important and widely discussed by both mathematicians and physicists. In [2], the concept of 2-Killing vector fields, as a new generalization of Killing vector fields, was first introduced and studied on Riemannian manifolds. In [3], Shenawy and Ünal studied 2-Killing vector fields on warped product manifolds, and applied their result to standard static space-times.

The definition of a semi-symmetric metric connection was given by H. Hayden in [4]. In 1970, K. Yano in [5] considered a semi-symmetric metric connection and studied some of its properties.

---

*Corresponding author.
E-mail addresses: quq453@nenu.edu.cn.
School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024, China

Motivated by the Yano's result, in [6], Sular and Özgür studied warped product manifolds with a semi-symmetric metric connection. In [7], professor Yong Wang studied multiply warped products with a semi-symmetric metric connection.

This paper is arranged as follows. In Section 2, we introduce a semi-symmetric metric connection, and then recall the definition of warped products and multiply warped products. In Section 3, we define a semi-symmetric metric Killing vector field, then study semi-symmetric metric Killing vector fields on warped products with a semi-symmetric metric connection. In Section 4, we study semi-symmetric metric Killing vector fields on multiply warped products with a semi-symmetric metric connection. In Section 5, we study Killing vector fields on multiply warped products. Finally in Section 6, we study 2-Killing vector fields on multiply warped products.

2 Preliminaries

Let M be a Riemannian manifold with Riemannian metric g. A linear connection $\overline{\nabla}$ on a Riemannian manifold M is called a semi-symmetric connection if the torsion tensor T of the connection $\overline{\nabla}$

$$T(X,Y)=\overline{\nabla}_XY-\overline{\nabla}_YX-[X,Y] \tag{1}$$

satisfies

$$T(X,Y)=\pi(Y)X-\pi(X)Y \tag{2}$$

where π is a 1-form associated with the vector field P on M defined by $\pi(X)=g(X,P)$. $\overline{\nabla}$ is called a semi-symmetric metric connection if it satisfies $\overline{\nabla}g=0$.

If ∇ is the Levi-Civita connection of M, the semi-symmetric metric connection $\overline{\nabla}$ is given by

$$\overline{\nabla}_XY=\nabla_XY+\pi(Y)X-g(X,Y)P, \tag{3}$$

(see [5]).

The $(1,3)$ type curvature R is defined by $R(X,Y)=\nabla_X\nabla_Y-\nabla_Y\nabla_X-\nabla_{[X,Y]}$, and $(0,4)$ type curvature is defined by $R(X,Y,Z,W)=g(R(X,Y)Z,W)$.

We recall the definition of warped product and multiply warped product as follows:

Definition 2.1. *Let* (M_1,g_1) *and* (M_2,g_2) *be two pseudo-Riemannian manifolds,* $f:M_1\to(0,\infty)$ *be a smooth function. The* ***warped product*** $M=M_1\times_f M_2$ *is the product manifold* $M_1\times M_2$ *with the metric tensor* $g=g_1\oplus f^2g_2$*. The function* f *is called the warping function of the warped product.*

Definition 2.2. *Let* (B,g_B) *and* (M_i,g_i) *be pseudo-Riemannian manifolds,* $f_i:B\to(0,\infty)$ *be smooth functions, where for each* $i\in\{1,\cdots,m\}$*. The* ***multiply warped product*** $M=B\times_{f_1}M_1\times_{f_2}M_2\times\cdots\times_{f_m}M_m$ *is the product manifold* $B\times M_1\times M_2\times\cdots\times M_m$ *with the metric tensor* $g=g_B\oplus f_1^2g_1\oplus f_2^2g_2\oplus\cdots\oplus f_m^2g_m$*. The function* f_i *is called the warping function of the multiply warped product.*

3 Semi-symmetric Metric Killing Vector Fields on Warped Products with a Semi-symmetric Metric Connection

Lemma 3.1. [6] *Let* $M=M_1\times_f M_2$ *be a warped product with* $\overline{\nabla}$*. If* $X_1,Y_1\in\Gamma(TM_1)$*,* $X_2,Y_2\in\Gamma(TM_2)$ *and* $P\in\Gamma(TM_1)$*, then*

$(1)\overline{\nabla}_{X_1}Y_1=\overline{\nabla}^1_{X_1}Y_1;$
$(2)\overline{\nabla}_{X_1}Y_2=\frac{X_1f}{f}Y_2;$
$(3)\overline{\nabla}_{Y_2}X_1=\left[\frac{X_1f}{f}+\pi(X_1)\right]Y_2;$
$(4)\overline{\nabla}_{X_2}Y_2=-fg_2(X_2,Y_2)\mathrm{grad}f+\nabla^2_{X_2}Y_2-f^2g_2(X_2,Y_2)P.$

Lemma 3.2. [6] *Let $M=M_1\times_f M_2$ be a warped product with $\overline{\nabla}$. If $X_1,Y_1\in\Gamma(TM_1)$, $X_2,Y_2\in\Gamma(TM_2)$ and $P\in\Gamma(TM_2)$, then*
$(1)\overline{\nabla}_{X_1}Y_1=\nabla^1_{X_1}Y_1-g_1(X_1,Y_1)P;$
$(2)\overline{\nabla}_{X_1}Y_2=\frac{X_1f}{f}Y_2+\pi(Y_2)X_1;$
$(3)\overline{\nabla}_{Y_2}X_1=\frac{X_1f}{f}Y_2;$
$(4)\overline{\nabla}_{X_2}Y_2=-fg_2(X_2,Y_2)\mathrm{grad}f+\nabla^2_{X_2}Y_2+\pi(Y_2)X_2-f^2g_2(X_2,Y_2)P.$

Lemma 3.3. *Let M be a pseudo-Riemannian manifold, $\zeta\in\Gamma(TM)$, $\mathcal{L}$ is the Lie derivative on M with respect to ζ, then:*

$$(\mathcal{L}_\zeta g)(X,Y)=g(\nabla_X\zeta,Y)+g(\nabla_Y\zeta,X)\tag{4}$$

for any vector fields $X,Y\in\Gamma(TM)$.

Definition 3.4. *Let (M,g) be a pseudo-Riemannian manifold, $\zeta\in\Gamma(TM)$, $\mathcal{L}$ is the Lie derivative on M with respect to ζ, then ζ is called **a Killing vector field** if*

$$\mathcal{L}_\zeta g=0\tag{5}$$

Now we define the semi-symmetric metric Lie derivative as follows:

Definition 3.5. *Let M be a pseudo-Riemannian manifold, $\zeta\in\Gamma(TM)$, $\overline{\mathcal{L}}$ is called **semi-symmetric metric Lie derivative** on M with respect to ζ, if it satisfied:*

$$(\overline{\mathcal{L}}_\zeta g)(X,Y)=g(\overline{\nabla}_X\zeta,Y)+g(\overline{\nabla}_Y\zeta,X)\tag{6}$$

for any vector fields $X,Y\in\Gamma(TM)$.

Definition 3.6. *Let (M,g) be a pseudo-Riemannian manifold, $\zeta\in\Gamma(TM)$, $\overline{\mathcal{L}}$ is the semi-symmetric metric Lie derivative on M with respect to ζ, then ζ is called **a semi-symmetric metric Killing vector field** if*

$$\overline{\mathcal{L}}_\zeta g=0\tag{7}$$

Using the symmetry in the equation (4), we can get:

Lemma 3.7. *If (M,g,∇) is a pseudo-Riemannian manifold, then $\zeta\in\Gamma(TM)$ is a Killing vector field if and only if:*

$$g(\nabla_X\zeta,X)=0\tag{8}$$

Using the symmetry in the equation (6), we can get:

Lemma 3.8. *If $(M,g,\overline{\nabla})$ is a pseudo-Riemannian manifold, then $\zeta\in\Gamma(TM)$ is a semi-symmetric metric Killing vector field if and only if:*

$$g(\overline{\nabla}_X\zeta,X)=0\tag{9}$$

Remark 3.9. *The relationship between semi-symmetric metric Killing vector field and Killing vector field:*

$$g(\overline{\nabla}_X\zeta, X) = g\big(\nabla_X\zeta + \pi(\zeta)X - g(X,\zeta)P, X\big) = g(\nabla_X\zeta, X) + \pi(\zeta)\|X\|^2 - \pi(X)g(X,\zeta).$$

So when $\pi(\zeta)\|X\|^2 = \pi(X)g(X,\zeta)$, *we have* $g(\overline{\nabla}_X\zeta, X) = 0 \Leftrightarrow g(\nabla_X\zeta, X) = 0$. *So we get:*

Proposition 3.10. *Let* $(M, g, \overline{\nabla})$ *be a pseudo-Riemannian manifold,* $\zeta \in \Gamma(TM)$, *if* $\pi(\zeta)\|X\|^2 = \pi(X)g(X,\zeta)$, *then the following two conditions are equivalent:*
(1) ζ *is a semi-symmetric metric Killing vector field on* M.
(2) ζ *is a Killing vector field on* M.
In particular, if $P = 0$, *then the above two conditions are equivalent.*

Remark 3.11. *It is easy to see that when* $P = 0$, *the semi-symmetric metric Lie derivative becomes to Lie derivative, and the semi-symmetric metric Killing vector field becomes to Killing vector field.*

Example 3.12. *The semi-symmetric metric Killing vector field on* I, *where* I *is an open interval in* $\mathbb{R}$, *and* $g_I = \mathrm{d}t^2$. *Suppose* $\zeta, X \in \Gamma(TI)$, *then we have* $\zeta = u\partial t$, $X = v\partial t$, *where* $u \neq 0$, $v \neq 0$, *then:*

$$\pi(\zeta)\|X\|^2 = g(P,\zeta)g(X,X) = g(P,u\partial t)g(v\partial t, v\partial t) = uv^2 g(P,\partial t),$$

$$\pi(X)g(X,\zeta) = g(P,X)g(X,\zeta) = g(P,v\partial t)g(u\partial t, v\partial t) = uv^2 g(P,\partial t),$$

which means $\pi(\zeta)\|X\|^2 = \pi(X)g(X,\zeta)$, *so by Proposition 3.10, we have the semi-symmetric metric Killing vector field on* I *is equivalent to the Killing vector field on* I, *and*

$$g(\overline{\nabla}_X\zeta, X) = g(\nabla_X\zeta, X) = g(\nabla_{v\partial t}u\partial t, v\partial t) = g(v\nabla_{\partial t}u\partial t, v\partial t) = g(v\dot{u}\partial t, v\partial t) = \dot{u}v^2 = 0,$$

since $v \neq 0$, *we have* $\dot{u} = 0$, *so* $u = a \in \mathbb{R} \setminus \{0\}$.

In a word, the semi-symmetric metric Killing vector field on I *and the Killing vector field on* I *has the form* $a\partial t$, *where* $a \in \mathbb{R} \setminus \{0\}$.

By Lemma 3.1 and straightforward computation, we can get:

Proposition 3.13. *Let* $M = M_1 \times_f M_2$ *be a warped product with* $\overline{\nabla}$, $\zeta \in \Gamma(TM)$, $P \in \Gamma(TM_1)$. *Then for any vector fields* $X, Y \in \Gamma(TM)$, *we have:*

$$\begin{aligned}(\overline{\mathcal{L}}_\zeta g)(X,Y) \;=\;& (\overline{\mathcal{L}}^1_{\zeta_1} g_1)(X_1,Y_1) + f^2(\mathcal{L}^2_{\zeta_2} g_1)(X_2,Y_2) + 2[f\zeta_1(f) + f^2\pi(\zeta_1)]g_2(X_2,Y_2)\\ &- f^2 g_2(X_2,\zeta_2)\pi(Y_1) - f^2 g_2(Y_2,\zeta_2)\pi(X_1)\end{aligned} \tag{10}$$

By Lemma 3.2 and straightforward computation, we can get:

Proposition 3.14. *Let* $M = M_1 \times_f M_2$ *be a warped product with* $\overline{\nabla}$, $\zeta \in \Gamma(TM)$, $P \in \Gamma(TM_2)$. *Then for any vector fields* $X, Y \in \Gamma(TM)$, *we have:*

$$\begin{aligned}(\overline{\mathcal{L}}_\zeta g)(X,Y) \;=\;& (\mathcal{L}^1_{\zeta_1} g_1)(X_1,Y_1) + f^2(\mathcal{L}^2_{\zeta_2} g_2)(X_2,Y_2) + 2\pi(\zeta_2)g(X,Y)\\ &+ 2f\zeta_1(f)g_2(X_2,Y_2) - \pi(Y_2)g(X,\zeta) - \pi(X_2)g(Y,\zeta)\end{aligned} \tag{11}$$

Using the equation (6) and Proposition 3.13, we have:

Corollary 3.15. *Let $M = M_1 \times_f M_2$ be a warped product with $\overline{\nabla}$, $\zeta \in \Gamma(TM)$, $P \in \Gamma(TM_1)$. Then for any vector field $X \in \Gamma(TM)$, we have:*

$$\begin{aligned} g(\overline{\nabla}_X \zeta, X) &= g_1(\overline{\nabla}^1_{X_1}\zeta_1, X_1) + f^2 g_2(\nabla^2_{X_2}\zeta_2, X_2) \\ &- f^2\pi(X_1)g_2(X_2,\zeta_2) + \left[f\zeta_1(f) + f^2\pi(\zeta_1)\right]\|X_2\|_2^2. \end{aligned} \tag{12}$$

Using the equation (6) and Proposition 3.14, we have:

Corollary 3.16. *Let $M = M_1 \times_f M_2$ be a warped product with $\overline{\nabla}$, $\zeta \in \Gamma(TM)$, $P \in \Gamma(TM_2)$. Then for any vector field $X \in \Gamma(TM)$, we have:*

$$\begin{aligned} g(\overline{\nabla}_X \zeta, X) &= g_1(\nabla^1_{X_1}\zeta_1, X_1) + f^2 g_2(\nabla^2_{X_2}\zeta_2, X_2) + \pi(\zeta_2)\|X\|^2 \\ &+ f\zeta_1(f)\|X_2\|_2^2 - \pi(X_2)g(X,\zeta). \end{aligned} \tag{13}$$

By Corollary 3.15, we can easily get:

Proposition 3.17. *Let $M = M_1 \times_f M_2$ be a warped product with $\overline{\nabla}$, $P \in \Gamma(TM_1)$, $\zeta = \zeta_1 + \zeta_2 \in \Gamma(TM)$. Then ζ is a semi-symmetric metric Killing vector field if one of the following conditions holds:*

(1) $\zeta = \zeta_1$, ζ_1 *is a semi-symmetric metric Killing vector field, and* $\zeta_1(f) + f\pi(\zeta_1) = 0$.

(2) $\zeta = \zeta_2$, ζ_2 *is a Killing vector field and* $\pi(X_1)g_2(X_2,\zeta_2) = 0$.

(3) $\zeta = \zeta_1 + \zeta_2$, ζ_1 *is a semi-symmetric metric Killing vector field,* ζ_2 *is a Killing vector field, and* $\zeta_1(f) + f\pi(\zeta_1) = 0$, $\pi(X_1)g_2(X_2,\zeta_2) = 0$.

Proposition 3.18. *Let $M = M_1 \times_f M_2$ be a warped product with $\overline{\nabla}$, $P \in \Gamma(TM_1)$, $\zeta = \zeta_1 + \zeta_2$ is a semi-symmetric metric Killing vector field, then:*

(1) ζ_1 *is a semi-symmetric metric Killing vector field on* M_1.

(2) ζ_2 *is a Killing vector field on* M_2 *if* $\zeta_1(f) + f\pi(\zeta_1) = 0$.

Now we recall the definition of generalized Robertson-Walker space-times as follow:

Definition 3.19. *A* ***generalized Robertson-Walker space-time*** *(M, g) is a warped product of the form $M = I \times_f M_2$ with the metric tensor $g = -\mathrm{d}t^2 \oplus f^2 g_2$, where $f : I \to (0,\infty)$ is smooth, I is an open interval in $\mathbb{R}$.*

Proposition 3.20. *Let $M = I \times_f M_2$ be a generalized Robertson-Walker space-time with $\overline{\nabla}$, $P = \partial t \in \Gamma(TI)$, $\zeta = a\partial t + \zeta_2 \in \Gamma(TM)$, $a \in \mathbb{R}\setminus\{0\}$. Then for any $X = u\partial t + X_2 \in \Gamma(TM)$, $u \in \mathbb{R}\setminus\{0\}$, we have: ζ is a semi-symmetric metric Killing vector field on M if ζ_2 is a Killing vector field on M_2 and $f = e^t$, $g_2(X_2,\zeta_2) = 0$.*

Proof. By Proposition 3.17.(3), we just need $a\partial t(f) + f\pi(a\partial t) = 0$, and $\pi(u\partial t)g_2(X_2,\zeta_2) = 0$.

On the one hand, $a\partial t(f) + f\pi(a\partial t) = a\dot{f} + fg(a\partial t, \partial t) = a\dot{f} - af = 0$. Consider that $a \neq 0$, we have $\dot{f} = f$, then $f = e^t$.

On the other hand, $\pi(u\partial t)g_2(X_2,\zeta_2) = g(u\partial t,\partial t)g_2(X_2,\zeta_2) = -ug_2(X_2,\zeta_2) = 0$, Consider that $u \neq 0$, we have $g_2(X_2,\zeta_2) = 0$. $\square$

By Corollary 3.16, we can easily get:

Proposition 3.21. *Let $M = M_1 \times_f M_2$ be a warped product with $\overline{\nabla}$, $P \in \Gamma(TM_2)$, $\zeta = \zeta_1 + \zeta_2 \in \Gamma(TM)$. Then ζ is a semi-symmetric metric Killing vector field if one of the following conditions holds:*
(1) $\zeta = \zeta_1$, *ζ_1 is a Killing vector field and* $f\zeta_1(f)\|X_2\|_2^2 - \pi(X_2)g_1(X_1, \zeta_1) = 0$.
(2) $\zeta = \zeta_2$, *ζ_2 is a Killing vector field and* $\pi(\zeta_2)\|X\|^2 - f^2\pi(X_2)g_2(X_2, \zeta_2) = 0$.
(3) $\zeta = \zeta_1 + \zeta_2$, *each ζ_i is a Killing vector field and* $f\zeta_1(f)\|X_2\|_2^2 + \pi(\zeta_2)\|X\|^2 - \pi(X_2)g(X, \zeta) = 0$.

Proposition 3.22. *Let $M = M_1 \times_f M_2$ be a warped product with $\overline{\nabla}$, $P \in \Gamma(TM_2)$, $\zeta = \zeta_1 + \zeta_2$ is a semi-symmetric metric Killing vector field, then:*
(1) *ζ_1 is a Killing vector field on M_1 if* $\pi(\zeta_2) = 0$.
(2) *ζ_2 is a Killing vector field on M_2 if* $\pi(\zeta_2) = 0$ *and* $f\zeta_1(f)\|X_2\|_2^2 - \pi(X_2)g(X, \zeta) = 0$.

Now we recall the definition of standard static space-times as follow:

Definition 3.23. *A **standard static space-time** (M, g) is a warped product of the form $M = M_1 \times_f I$ with the metric tensor $g = g_1 \oplus (-f^2)\mathrm{d}t^2$, where $f : M_1 \to (0, \infty)$ is smooth, I is an open interval in $\mathbb{R}$.*

Proposition 3.24. *Let $M = M_1 \times_f I$ be a standard static space-time with $\overline{\nabla}$, $P = \partial t \in \Gamma(TI)$, $\zeta = \zeta_1 + a\partial t \in \Gamma(TM)$, $a \in \mathbb{R} \setminus \{0\}$. Then for any $X = X_1 + u\partial t \in \Gamma(TM)$, $u \in \mathbb{R} \setminus \{0\}$, we have: ζ is a semi-symmetric metric Killing vector field on M if ζ_1 is a Killing vector field on M_1 and* $ufg_1(X_1, \zeta_1) - u^2\zeta_1(f) - af\|X_1\|_1^2 = 0$.

Proof. By Proposition 3.21.(3), we just need

$$f\zeta_1(f)\|u\partial t\|_2^2 + \pi(a\partial t)\|X_1 + u\partial t\|^2 - \pi(u\partial t)g(X_1 + u\partial t, \zeta_1 + a\partial t) = 0.$$

$$\Rightarrow f\zeta_1(f)g_2(u\partial t, u\partial t) + g(a\partial t, \partial t)\Big[\|X_1\|_1^2 + f^2 g_2(u\partial t, u\partial t)\Big] - g(u\partial t, \partial t)\Big[g_1(X_1, \zeta_1) + f^2 g_2(u\partial t, a\partial t)\Big] = 0,$$

$$\Rightarrow -u^2 f\zeta_1(f) - af^2\Big[\|X_1\|_1^2 - u^2 f^2\Big] + uf^2\Big[g_1(X_1, \zeta_1) - auf^2\Big] = 0,$$

$$\Rightarrow uf^2 g_1(X_1, \zeta_1) - u^2 f\zeta_1(f) - af^2\|X_1\|_1^2 = 0,$$

since $f > 0$, we get $ufg_1(X_1, \zeta_1) - u^2\zeta_1(f) - af\|X_1\|_1^2 = 0$. □

4 Semi-symmetric Metric Killing Vector Fields on Multiply Warped Products with a Semi-symmetric Metric Connection

Lemma 4.1. [7] *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product with $\overline{\nabla}$. If $X_B, Y_B \in \Gamma(TB)$, $X_i, Y_i \in \Gamma(TM_i)$ and $P \in \Gamma(TB)$, $i \in \{1, \cdots, m\}$, then*
(1) $\overline{\nabla}_{X_B}Y_B = \overline{\nabla}^B_{X_B}Y_B$;
(2) $\overline{\nabla}_{X_B}Y_i = \frac{X_B(f_i)}{f_i}Y_i$;
(3) $\overline{\nabla}_{Y_i}X_B = \Big[\frac{X_B(f_i)}{f_i} + \pi(X_B)\Big]Y_i$;
(4) $\overline{\nabla}_{X_i}Y_j = 0$, *if* $i \neq j$;
(5) $\overline{\nabla}_{X_i}Y_i = -f_i g_i(X_i, Y_i)\mathrm{grad}f_i + \nabla^i_{X_i}Y_i - f_i^2 g_i(X_i, Y_i)P$.

Lemma 4.2. [7] *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$*. If* $X_B, Y_B \in \Gamma(TB)$*,* $X_i, Y_i \in \Gamma(TM_i)$ *and* $P \in \Gamma(TM_r)$ *for a fixed* r*,* $i \in (1, \cdots, m)$*, then*
(1)$\overline{\nabla}_{X_B} Y_B = \nabla^B_{X_B} Y_B - g_B(X_B, Y_B)P$;
(2)$\overline{\nabla}_{X_B} Y_i = \frac{X_B(f_i)}{f_i} Y_i + \pi(Y_i) X_B$;
(3)$\overline{\nabla}_{Y_i} X_B = \frac{X_B(f_i)}{f_i} Y_i$;
(4)$\overline{\nabla}_{X_i} Y_j = \pi(Y_j) X_i$*, if* $i \neq j$;
(4)$\overline{\nabla}_{X_i} Y_i = -f_i g_i(X_i, Y_i)\mathrm{grad} f_i + \nabla^i_{X_i} Y_i + \pi(Y_i) X_i - f_i^2 g_i(X_i, Y_i)P$.

By Lemma 4.1 and straightforward computation, we can get:

Proposition 4.3. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$*,* $\zeta \in \Gamma(TM)$*,* $P \in \Gamma(TB)$*. Then for any vector fields* $X, Y \in \Gamma(TM)$*, we have:*

$$\begin{aligned}(\overline{\mathcal{L}}_\zeta g)(X, Y) &= (\overline{\mathcal{L}}^B_{\zeta_B} g_B)(X_B, Y_B) + \sum_{i=1}^m f_i^2 (\mathcal{L}^i_{\zeta_i} g_i)(X_i, Y_i) + \sum_{i=1}^m 2[f_i \zeta_B(f_i) + f_i^2 \pi(\zeta_B)] g_i(X_i, Y_i) \\ &\quad - \sum_{i=1}^m f_i^2 g_i(X_i, \zeta_i)\pi(Y_B) - \sum_{i=1}^m f_i^2 g_i(Y_i, \zeta_i)\pi(X_B) \qquad (14)\end{aligned}$$

By Lemma 4.2 and straightforward computation, we can get:

Proposition 4.4. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$*,* $\zeta \in \Gamma(TM)$*,* $P \in \Gamma(TM_r)$ *for a fixed* r*. Then for any vector fields* $X, Y \in \Gamma(TM)$*, we have:*

$$\begin{aligned}(\overline{\mathcal{L}}_\zeta g)(X, Y) &= (\mathcal{L}^B_{\zeta_B} g_B)(X_B, Y_B) + \sum_{i=1}^m f_i^2 (\mathcal{L}^i_{\zeta_i} g_i)(X_i, Y_i) + \sum_{i=1}^m 2\pi(\zeta_i) g(X, Y) \\ &+ \sum_{i=1}^m 2 f_i \zeta_B(f_i) g_i(X_i, Y_i) - \sum_{i=1}^m \pi(Y_i) g(X, \zeta) - \sum_{i=1}^m \pi(X_i) g(Y, \zeta) \qquad (15)\end{aligned}$$

Using the equation (6) and Proposition 4.3, we have:

Corollary 4.5. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$*,* $P \in \Gamma(TB)$*. Then for any vector field* $X \in \Gamma(TM)$*, we have:*

$$\begin{aligned}g(\overline{\nabla}_X \zeta, X) &= g_B(\overline{\nabla}^B_{X_B} \zeta_B, X_B) + \sum_{i=1}^m f_i^2 g_i(\nabla^i_{X_i} \zeta_i, X_i) \\ &- \sum_{i=1}^m f_i^2 \pi(X_B) g_i(X_i, \zeta_i) + \sum_{i=1}^m \left[f_i \zeta_B(f_i) + f_i^2 \pi(\zeta_B)\right] \|X_i\|_i^2. \qquad (16)\end{aligned}$$

Using the equation (6) and Proposition 4.4, we have:

Corollary 4.6. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$*,* $\zeta \in \Gamma(TM)$*,* $P \in \Gamma(TM_r)$ *for a fixed* r*. Then for any vector field* $X \in \Gamma(TM)$*, we have:*

$$\begin{aligned}g(\overline{\nabla}_X \zeta, X) &= g_B(\nabla^B_{X_B} \zeta_B, X_B) + \sum_{i=1}^m f_i^2 g_i(\nabla^i_{X_i} \zeta_i, X_i) + \sum_{i=1}^m \pi(\zeta_i)\|X\|^2 \\ &+ \sum_{i=1}^m f_i \zeta_B(f_i) \|X_i\|_i^2 - \sum_{i=1}^m \pi(X_i) g(X, \zeta). \qquad (17)\end{aligned}$$

By Corollary 4.5, we can easily get:

Proposition 4.7. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$, $P \in \Gamma(TB)$, $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$. *Then* ζ *is a semi-symmetric metric Killing vector field if one of the following conditions holds:*

(1) $\zeta = \zeta_B$, ζ_B *is a semi-symmetric metric Killing vector field, and* $\sum\limits_{i=1}^{m} \big[f_i\zeta_B(f_i) + f_i^2\pi(\zeta_B)\big]\|X_i\|_i^2 = 0.$

(2) $\zeta = \zeta_i$ *for a fixed* i, ζ_i *is a Killing vector field and* $\pi(X_B)g_i(X_i, \zeta_i) = 0$.

(3) $\zeta = \zeta_B + \zeta_i$ *for a fixed* i, ζ_B *is a semi-symmetric metric Killing vector field,* ζ_i *is a Killing vector field, and* $\big[\zeta_B(f_i) + f_i\pi(\zeta_B)\big]\|X_i\|_i^2 - f_i\pi(X_B)g_i(X_i, \zeta_i) = 0$.

(4) $\zeta = \sum\limits_{i=1}^{m} \zeta_i$, *each* ζ_i *is a Killing vector field and* $\sum\limits_{i=1}^{m} f_i^2\pi(X_B)g_i(X_i, \zeta_i) = 0$, *for any* $i \in \{1, \cdots, m\}$.

(5) $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i$, ζ_B *is a semi-symmetric metric Killing vector field, each* ζ_i *is a Killing vector field and* $\sum\limits_{i=1}^{m} \big[f_i\zeta_B(f_i) + f_i^2\pi(\zeta_B)\big]\|X_i\|_i^2 - \sum\limits_{i=1}^{m} f_i^2\pi(X_B)g_i(X_i, \zeta_i) = 0$, *for any* $i \in \{1, \cdots, m\}$.

Proposition 4.8. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$, $P \in \Gamma(TB)$, $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i$ *is a semi-symmetric metric Killing vector field. Then:*

(1) ζ_B *is a semi-symmetric metric Killing vector field on* B.

(2) *each* ζ_i *is a Killing vector field on* M_i *if* $\sum\limits_{i=1}^{m} \big[f_i\zeta_B(f_i) + f_i^2\pi(\zeta_B)\big]\|X_i\|_i^2 = 0$.

By Corollary 4.6, we can easily get:

Proposition 4.9. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$, $P \in \Gamma(TM_r)$ *for a fixed* r, $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$. *Then* ζ *is a semi-symmetric metric Killing vector field if one of the following conditions holds:*

(1) $\zeta = \zeta_B$, ζ_B *is a Killing vector field, and* $\sum\limits_{i=1}^{m} \big[f_i\zeta_B(f_i)\|X_i\|_i^2 - \pi(X_i)g_B(X_B, Y_B)\big] = 0$.

(2) $\zeta = \zeta_i$ *for a fixed* $i \neq r$, ζ_i *is a Killing vector field ;*
$\zeta = \zeta_i$ *for a fixed* $i = r$, ζ_i *is a Killing vector field and* $\pi(\zeta_i)\|X_i\|_i^2 - \pi(X_i)g(X_i, \zeta_i) = 0$.

(3) $\zeta = \zeta_B + \zeta_i$ *for a fixed* $i \neq r$, ζ_B *is a Killing vector field,* ζ_i *is a Killing vector field and* $\zeta_B(f_i) = 0$.
$\zeta = \zeta_B + \zeta_i$ *for a fixed* $i = r$, ζ_B *is a Killing vector field,* ζ_i *is a Killing vector field, and* $\zeta_B(f_i)\|X_i\|_i^2 + f_i\pi(\zeta_i)\|X_i\|_i^2 - f_i\pi(X_i)g_i(X_i, \zeta_i) = 0$.

(4) $\zeta = \sum\limits_{i=1}^{m} \zeta_i$, *each* ζ_i *is a Killing vector field, and* $\sum\limits_{i=1}^{m} f_i^2\big[\pi(\zeta_i)\|X_i\|_i^2 - \pi(X_i)g_i(X_i, \zeta_i)\big] = 0$, *for* $i \in \{1, \cdots, m\}$.

(5) $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i$, ζ_B *is a Killing vector field, each* ζ_i *is a Killing vector field, and*
$\sum\limits_{i=1}^{m} \big[f_i\zeta_B(f_i)\|X_i\|_i^2 + \pi(\zeta_i)\|X\|^2 - \pi(X_i)g(X, \zeta)\big] = 0$.

Proposition 4.10. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product with* $\overline{\nabla}$, $P \in \Gamma(TM_r)$ *for a fixed* r, $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i$ *is a semi-symmetric metric Killing vector field, then:*

(1) ζ_B *is a Killing vector field on* B *if* $\sum\limits_{i=1}^{m} \pi(\zeta_i) = 0$.

(2) *each* ζ_i *is a Killing vector field on* M_i *if* $\sum\limits_{i=1}^{m} \pi(\zeta_i) = 0$ *and* $\sum\limits_{i=1}^{m} \big[f_i\zeta_B(f_i)\|X_i\|_i^2 - \pi(X_i)g(X, \zeta)\big] = 0$.

5 Killing Vector Fields on Multiply Warped Products

Notice that when $P = 0$, the semi-symmetric metric Lie derivative becomes to Lie derivative, and the semi-symmetric metric Killing vector field becomes to Killing vector field. So the following four results are special cases of Section 4 when $P = 0$.

Proposition 5.1. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product, $\zeta \in \Gamma(TM)$. Then for any vector fields $X, Y \in \Gamma(TM)$, we have:*

$$(\mathcal{L}_\zeta g)(X,Y) = (\mathcal{L}^B_{\zeta_B} g_B)(X_B, Y_B) + \sum_{i=1}^m f_i^2 (\mathcal{L}^i_{\zeta_i} g_i)(X_i, Y_i) + 2\sum_{i=1}^m f_i \zeta_B(f_i) g_i(X_i, Y_i). \tag{18}$$

Corollary 5.2. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product, $\zeta \in \Gamma(TM)$. Then for any vector field $X \in \Gamma(TM)$, we have:*

$$g(\nabla_X \zeta, X) = g_B(\nabla^B_{X_B} \zeta_B, X_B) + \sum_{i=1}^m f_i^2 g_i(\nabla^i_{X_i} \zeta_i, X_i) + \sum_{i=1}^m f_i \zeta_B(f_i) \|X_i\|_i^2. \tag{19}$$

Proposition 5.3. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product, $\zeta = \zeta_B + \sum_{i=1}^m \zeta_i \in \Gamma(TM)$. Then ζ is a Killing vector field if one of the following conditions holds:*

(1) *$\zeta = \zeta_B$, ζ_B is a Killing vector field, and $\sum_{i=1}^m f_i \zeta_B(f_i) \|X_i\|_i^2 = 0$.*

(2) *$\zeta = \zeta_i$ for a fixed i, ζ_i is a Killing vector field.*

(3) *$\zeta = \zeta_B + \zeta_i$ for a fixed i, ζ_B is a Killing vector field, ζ_i is a Killing vector field, and $\zeta_B(f_i) = 0$.*

(4) *$\zeta = \sum_{i=1}^m \zeta_i$, each ζ_i is a Killing vector field.*

(5) *$\zeta = \zeta_B + \sum_{i=1}^m \zeta_i$, ζ_B is a Killing vector field, each ζ_i is a Killing vector field, $\sum_{i=1}^m f_i \zeta_B(f_i) \|X_i\|_i^2 = 0$.*

Proposition 5.4. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product, $\zeta = \zeta_B + \sum_{i=1}^m \zeta_i$ is a Killing vector field. Then:*

(1) *ζ_B is a Killing vector field on B.*

(2) *each ζ_i is a Killing vector field on M_i if $\sum_{i=1}^m f_i \zeta_B(f_i) \|X_i\|_i^2 = 0$, for $i \in \{1, \cdots, m\}$.*

6 2-Killing Vector Fields on Multiply Warped Products

Definition 6.1. [2] *Let M be a pseudo-Riemannian manifold, $\zeta \in \Gamma(TM)$ is called a* ***2-Killing vector field****, if*

$$\mathcal{L}_\zeta \mathcal{L}_\zeta g = 0 \tag{20}$$

where $\mathcal{L}_\zeta$ is the Lie derivative in the direction of ζ on M.

Proposition 6.2. [2] *Let M be a pseudo-Riemannian manifold, $\zeta \in \Gamma(TM)$, then:*

$$(\mathcal{L}_\zeta \mathcal{L}_\zeta g)(X,Y) = g(\nabla_\zeta \nabla_X \zeta - \nabla_{[\zeta,X]}\zeta, Y) + g(X, \nabla_\zeta \nabla_Y \zeta - \nabla_{[\zeta,Y]}\zeta) + 2g(\nabla_X \zeta, \nabla_Y \zeta) \tag{21}$$

for any vector fields $X, Y \in \Gamma(TM)$.

Corollary 6.3. [2] *Let M be a pseudo-Riemannian manifold, $\zeta \in \Gamma(TM)$, then ζ is a 2-Killing vector field if and only if:*

$$R(\zeta, X, X, \zeta) = g(\nabla_X \zeta, \nabla_X \zeta) + g(\nabla_X \nabla_\zeta \zeta, X) \tag{22}$$

for any vector field $X \in \Gamma(TM)$.

Lemma 6.4. [8] *Let M be a pseudo-Riemannian manifold, $\zeta \in \Gamma(TM)$ is a Killing vector field of constant length, then $\nabla_\zeta \zeta = 0$.*

Then by Corollary 6.3 and Lemma 6.4, we have:

Corollary 6.5. *Let M be a pseudo-Riemannian manifold, $\zeta \in \Gamma(TM)$, ζ is a Killing vector field of constant length and also a 2-Killing vector field, then:*

$$R(\zeta, X, X, \zeta) = g(\nabla_X \zeta, \nabla_X \zeta) \geq 0 \tag{23}$$

for any vector field $X \in \Gamma(TM)$.

Lemma 6.6. [2] *Let ζ be a 2-Killing vector field on the compact $n-$dimensional pseudo-Riemannian manifold (M, g) without boundary. If $Ric(\zeta, \zeta) \leq 0$, then ζ is a parallel vector field and*

$$Tr\Big(g(\nabla \zeta, \nabla \zeta)\Big) = 0. \tag{24}$$

Lemma 6.7. [9] *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product. If $X_B, Y_B \in \Gamma(TB)$, $X_i, Y_i \in \Gamma(TM_i)$, $i \in \{1, \cdots, m\}$, then*
(1)$\nabla_{X_B} Y_B = \nabla^B_{X_B} Y_B$;
(2)$\nabla_{X_B} Y_i = \nabla_{Y_i} X_B = \frac{X_B(f_i)}{f_i} Y_i$;
(4)$\nabla_{X_i} Y_j = 0$, *if $i \neq j$;*
(4)$\nabla_{X_i} Y_i = \nabla^i_{X_i} Y_i - f_i g_i(X_i, Y_i) \mathrm{grad} f_i$.

By Lemma 6.7 and straightforward computation, we can get:

Proposition 6.8. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product. $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$, then we have:*

$$\begin{aligned}(\mathcal{L}_\zeta \mathcal{L}_\zeta g)(X, Y) &= (\mathcal{L}^B_{\zeta_B} \mathcal{L}^B_{\zeta_B} g_B)(X_B, Y_B) + \sum_{i=1}^{m} f_i^2 (\mathcal{L}^i_{\zeta_i} \mathcal{L}^i_{\zeta_i} g_i)(X_i, Y_i) + 4 \sum_{i=1}^{m} f_i \zeta_B(f_i) (\mathcal{L}^i_{\zeta_i} g_i)(X_i, Y_i) \\ &+ 2 \sum_{i=1}^{m} f_i \zeta_B\Big(\zeta_B(f_i)\Big) g_i(X_i, Y_i) + 2 \sum_{i=1}^{m} \zeta_B(f_i) \zeta_B(f_i) g_i(X_i, Y_i)\end{aligned} \tag{25}$$

By Proposition 6.8, we have:

Corollary 6.9. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product. $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$, if ζ is a 2-Killing vector field, then:*
(1) *ζ_B is a 2-Killing vector field.*
(2) *each ζ_i is a 2-Killing vector field if $\zeta_B(f_i) = 0$, for $i \in \{1, \cdots, m\}$.*

Corollary 6.10. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product. $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$, if ζ_B and each ζ_i are 2-Killing vector fields on B and M_i respectively, then ζ is a 2-Killing vector field on M if and only if one of the following conditions holds:*
(1) $\zeta_B(f_i) = 0$.
(2) *each ζ_i is a homothetic vector field on M_i with homothetic factor c_i(i.e, $\mathcal{L}^i_{\zeta_i} g_i = c_i g_i$) such that*

$$f_i \zeta_B\Big(\zeta_B(f_i)\Big) + \zeta_B(f_i)\zeta_B(f_i) = -2c_i f_i \zeta_B(f_i). \tag{26}$$

for any $i \in \{1, \cdots, m\}$.

Corollary 6.11. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product. $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$, then ζ is a 2-Killing vector field on M if one of the following conditions holds:*
(1) *ζ_B and each ζ_i are 2-Killing vector fields on B and M_i respectively, and $\zeta_B(f_i) = 0$, for any $i \in \{1, \cdots, m\}$.*
(2) *$\zeta = \sum\limits_{i=1}^{m} \zeta_i$ and ζ_i is a 2-Killing vector field on M_i, for any $i \in \{1, \cdots, m\}$.*

Proposition 6.12. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product, $\dim B = n$, $\dim M_i = n_i$, $i \in \{1, \cdots, m\}$, $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$, then*

$$\begin{aligned} Tr\Big(g(\nabla\zeta, \nabla\zeta)\Big) &= Tr\Big(g_B(\nabla^B\zeta_B, \nabla^B\zeta_B)\Big) + \sum_{i=1}^{m} Tr\Big(g_i(\nabla^i\zeta_i, \nabla^i\zeta_i)\Big) + 2\sum_{i=1}^{m} \|\zeta_i\|_i^2 \|\mathrm{grad} f_i\|_B^2 \\ &+ \sum_{i=1}^{m} \frac{n_i}{f_i^2}\Big(\zeta_B(f_i)\Big)^2 + 2\sum_{i=1}^{m} \frac{\zeta_B(f_i)}{f_i}\mathrm{div}_i\zeta_i. \end{aligned} \tag{27}$$

Proof. Suppose $\{e_1^B, e_2^B, \cdots, e_n^B\}$ is an orthonormal frame of B, $\{e_1^i, e_2^i, \cdots, e_{n_i}^i\}$ is an orthonormal frame of M_i, $i \in \{1, \cdots, m\}$. Then $\{e_1^B, e_2^B, \cdots, e_n^B, e_1^1, e_2^1, \cdots, e_{n_1}^1, \cdots, e_1^m, e_2^m, \cdots, e_{n_m}^m\}$ is an orthonormal frame of M. Then for any $\zeta \in \Gamma(TM)$, we have:

$$Tr\Big(g(\nabla\zeta, \nabla\zeta)\Big) = \sum_{i=1}^{n} g(\nabla_{e_i^B}\zeta, \nabla_{e_i^B}\zeta) + \frac{1}{f_1^2}\sum_{i=1}^{n_1} g(\nabla_{e_i^1}\zeta, \nabla_{e_i^1}\zeta) + \cdots + \frac{1}{f_m^2}\sum_{i=1}^{n_m} g(\nabla_{e_i^m}\zeta, \nabla_{e_i^m}\zeta).$$

By Lemma 6.7 and straightforward computation, we get

$$\sum_{i=1}^{n} g(\nabla_{e_i^B}\zeta, \nabla_{e_i^B}\zeta) = Tr\Big(g_B(\nabla^B\zeta_B, \nabla^B\zeta_B)\Big) + \sum_{i=1}^{m} \|\zeta_i\|_i^2 \|\mathrm{grad} f_i\|_B^2,$$

$$\frac{1}{f_i^2}\sum_{j=1}^{n_i} g(\nabla_{e_j^i}\zeta, \nabla_{e_j^i}\zeta) = \frac{n_i}{f_i^2}\Big(\zeta_B(f_i)\Big)^2 + Tr\Big(g_i(\nabla^i\zeta_i, \nabla^i\zeta_i)\Big) + \|\zeta_i\|_i^2\|\mathrm{grad} f_i\|_B^2 + 2\frac{\zeta_B(f_i)}{f_i}\mathrm{div}_i\zeta_i.$$

So we get the equation (27). □

Theorem 6.13. *Let $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ be a multiply warped product. $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i \in \Gamma(TM)$, B is a compact manifold without boundary, each M_i is a compact manifold without boundary, then:*

(1) $\zeta = \zeta_B + \sum\limits_{i=1}^{m} \zeta_i$ *is parallel if* ζ_B *is a 2-Killing vector field, each* ζ_i *is a 2-Killing vector field,* $Ric^B(\zeta_B, \zeta_B) \leq 0$, $Ric^i(\zeta_i, \zeta_i) \leq 0$, *and* f_i *is constant, for* $i \in \{1, \cdots, m\}$.

(2) $\zeta = \zeta_B$ *is parallel if* ζ_B *is a 2-Killing vector field,* $Ric^B(\zeta_B, \zeta_B) \leq 0$, *and* $\zeta_B(f_i) = 0$.

(3) $\zeta = \zeta_B + \zeta_i$ *is parallel for a fixed* i *if* ζ_B *is a 2-Killing vector field,* ζ_i *is a 2-Killing vector field,* $Ric^B(\zeta_B, \zeta_B) \leq 0$, $Ric^i(\zeta_i, \zeta_i) \leq 0$, *and* f_i *is constant.*

(4) $\zeta = \zeta_i$ *is parallel for a fixed* i *if* ζ_i *is a 2-Killing vector field,* $Ric^i(\zeta_i, \zeta_i) \leq 0$, *and* f_i *is constant.*

(5) $\zeta = \sum\limits_{i=1}^{m} \zeta_i$ *is parallel if each* ζ_i *is a 2-Killing vector field,* $Ric^i(\zeta_i, \zeta_i) \leq 0$, *and* f_i *is constant, for* $i \in \{1, \cdots, m\}$.

Proof. (1) Since ζ_B is a 2-Killing vector field, each ζ_i is a 2-Killing vector field, $Ric^B(\zeta_B, \zeta_B) \leq 0$, $Ric^i(\zeta_i, \zeta_i) \leq 0$, B is a compact manifold without boundary, each M_i is a compact manifold without boundary, by Lemma 6.6, we have $Tr\Big(g_B(\nabla^B \zeta_B, \nabla^B \zeta_B)\Big) = 0$, $Tr\Big(g_i(\nabla^i \zeta_i, \nabla^i \zeta_i)\Big) = 0$. Then for constant function f_i, using Proposition 6.12, we get :

$$Tr\Big(g(\nabla\zeta, \nabla\zeta)\Big) = 0.$$

Thus ζ is a parallel vector field with respect to the metric g.

Using the same proof method, we can easily get (2), (3), (4), (5). □

Theorem 6.14. *Let* $M = B \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product.* ζ *is a non-trivial 2-Killing vector field on* M, *let* K *denote the sectional curvature, then:*
(1) *If* $\nabla_\zeta \zeta$ *is parallel along a curve* γ, *then* $K(\zeta, \dot{\gamma}) \geq 0$.
(2) *If* ζ *is a Killing vector field of constant length, then* $K(\zeta, \dot{\gamma}) \geq 0$.

Proof. (1) By Corollary 6.3, we have $R(\zeta, X, X, \zeta) = g(\nabla_X \zeta, \nabla_X \zeta) + g(\nabla_X \nabla_\zeta \zeta, X)$. Take $X = \dot{\gamma}$, since $\nabla_\zeta \zeta$ is parallel along a curve γ, then we have $\nabla_X \nabla_\zeta \zeta = 0$, so

$$R(\zeta, X, X, \zeta) = g(\nabla_X \zeta, \nabla_X \zeta) = \|\nabla_X \zeta\| \geq 0$$

$$\Rightarrow -R(\zeta, X, \zeta, X) = \|\nabla_X \zeta\| \geq 0$$

$$\Rightarrow K(\zeta, \dot{\gamma}) = -\frac{R(\zeta, X, \zeta, X)}{A^2(\zeta, \dot{\gamma})} \geq 0,$$

where $A(\zeta, \dot{\gamma})$ is area of the parallelogram generated by ζ and $\dot{\gamma}$.
(2) Take $X = \dot{\gamma}$, by Corollary 6.5, we have

$$R(\zeta, X, X, \zeta) = g(\nabla_X \zeta, \nabla_X \zeta) \geq 0$$

$$\Rightarrow -R(\zeta, X, \zeta, X) = \|\nabla_X \zeta\| \geq 0$$

$$\Rightarrow K(\zeta, \dot{\gamma}) = -\frac{R(\zeta, X, \zeta, X)}{A^2(\zeta, \dot{\gamma})} \geq 0,$$

where $A(\zeta, \dot{\gamma})$ is area of the parallelogram generated by ζ and $\dot{\gamma}$. □

From [2], we can see that $(at - b)^{\frac{1}{3}} \partial t$ is a 2-Killing vector field on I, where $a, b \in \mathbb{R}$, $t \in I$, and $t \neq \frac{b}{a}$. Then by Corollary 6.10, we have:

Proposition 6.15. *Let* $M = I \times_{f_1} M_1 \times_{f_2} M_2 \times \cdots \times_{f_m} M_m$ *be a multiply warped product.* $\zeta = (at-b)^{\frac{1}{3}}\partial t + \sum_{i=1}^{m} \zeta_i \in \Gamma(TM)$. *Suppose each* ζ_i *is a 2-Killing vector field on* M_i*, then* ζ *is a 2-Killing vector field on* M *if each* ζ_i *is a homothetic vector field on* M_i *with* c_i *satisfying:*

$$\frac{a}{3} f_i \dot{f}_i + (f_i \ddot{f}_i + \dot{f_i}^2)(at-b) = -2c_i f_i \dot{f}_i (at-b)^{\frac{2}{3}}. \tag{28}$$

We recall the definition of generalized Kasner space-times in [9].

Definition 6.16. *A* ***generalized Kasner space-time*** (M, g) *is a Lorentzian multiply warped product of the form* $M = I \times_{\phi^{p_1}} M_1 \times_{\phi^{p_2}} M_2 \times \cdots \times_{\phi^{p_m}} M_m$ *with the metric tensor* $g = -dt^2 \oplus \phi^{2p_1} g_{M_1} \oplus \phi^{2p_2} g_{M_2} \oplus \cdots \oplus \phi^{2p_m} g_{M_m}$*, where* $\phi : I \to (0, \infty)$ *is smooth and* $p_i \in \mathbb{R}$*, for any* $i \in \{1, \ldots, m\}$ *and also* $I = (t_1, t_2)$*.*

Then consider that $f_i = \phi^{p_i}$, by Proposition 6.15, we get:

Proposition 6.17. *Let* $M = I \times_{\phi^{p_1}} M_1 \times_{\phi^{p_2}} M_2 \times \cdots \times_{\phi^{p_m}} M_m$ *be a generalized Kasner space-time.* $\zeta = (at-b)^{\frac{1}{3}}\partial t + \sum_{i=1}^{m} \zeta_i \in \Gamma(TM)$. *Suppose each* ζ_i *is a 2-Killing vector field on* M_i*, then* ζ *is a 2-Killing vector field on* M *if each* ζ_i *is a homothetic vector field on* M_i *with* c_i *satisfying:*

$$\frac{a}{3} + \frac{2p_i - 1}{\phi}(at-b) = -2c_i (at-b)^{\frac{2}{3}}. \tag{29}$$

Acknowledgement. We would like to thank the referee for his(her) careful reading and helpful comments.

References

[1] R. Bishop, B. O'Neill, Manifolds of negative curvature, Trans. Am. Math. Soc. 145(1969) 1-49.

[2] T. oprea, 2-Killing vector fields on Riemannian manifolds, Balkan Journal of Geometry and Its Applications, 13(2008), No.1, 87-92.

[3] S. Shenawy, B. Ünal, 2-Killing vector fields on standard static space-times, arXiv:1411.6207.

[4] H. A. Hayden, Subspace of a space with torsion, Proc. Lond. Math. Soc. 34(1932) 27-50.

[5] K. Yano, On semi-symmetric metric connection, Rev. Roumaine Math. Pures Appl. 15(1970) 1579-1586.

[6] S. Sular and C. Özgür, Warped products with a semi-symmetric metric connection, Taiwanese J. Math. 15 (2011), No. 4, 1701-1719.

[7] Y. Wang, Multiply warped products with a semi-symmetric metric connection, Abstract and Applied Analysis. Vol 2014, Article ID 742371, 12pages.

[8] Yu. G. Nikonorov, Killing vector fields of constant length on compact homogeneous Riemannian manifolds, arXiv:1504.03432.

[9] F. Dobarro, B. Ünal, Curvature of multiply warped products, J. Geom. Phys. 55(2005) 75-106.